

URO-T

To 100 mL URO medium add 50 mg Tris(hydroxymethyl) aminomethane.

URO

NH ₄ NO ₃	0.5 mg
β-Na ₂ glycerophosphate · 5H ₂ O	0.4 mg
MgSO ₄ · 7H ₂ O	1 mg
CaCl ₂ · 2H ₂ O	1 mg
KCl	0.1 mg
Thiamine HCl	1 μg
Vitamin B ₁₂	0.01 μg
Biotin	0.01 μg
Fe-EDTA	0.05 mg
PIV metals	0.1 mL
Distilled water	99.9 mL
pH 7.5 ¹⁾	

1) pH is adjusted to 7.5 with 0.1 mol/L HCl.

Reference

- Kimura, B., Ishida, Y. 1985 Photophagotrophy in *Uroglena americana*, Chrysophyceae. *Jpn. J. Limnol.*, **46**, 315-318.
- Nakahara, H., Sako, Y. 1987 2. Life history of freshwater phytoplankton [2. Tansui syokubutsu purankuton no seikatsushi]. In *Freshwater red tide [Tansui Akashio]*, Ed. by Kadota, H., Kôseisya-Kôseikaku, Tokyo, p. 21-77 (in Japanese without English title).

P IV metals

Na ₂ EDTA · 2H ₂ O	100 mg
FeCl ₃ · 6H ₂ O	19.6 mg
MnCl ₂ · 4H ₂ O	3.6 mg
ZnCl ₂ ¹⁾	1.04 mg
CoCl ₂ · 6H ₂ O	0.4 mg
Na ₂ MoO ₄ · 2H ₂ O	0.25 mg
Distilled water	100 mL

1) In the NIES-Collection, 1.04 mg ZnCl₂ is replaced by 2.2mg ZnSO₄ · 7H₂O.

Reference

- Provasoli, L., Pintner, I. J. 1959 Artificial media for fresh-water algae: problems and suggestions. In *The Ecology of Algae. Spec. Pub. No. 2.*, Eds. by Tryon, C. A., Jr. & Hartmann, R. T., Pymatuning Laboratory of Field Biology, University of Pittsburgh, Pittsburgh, p. 84-96.